

Vaccines with Diluents: How to Use Them

Be sure to reconstitute the following vaccines correctly before administering them! Reconstitution means that the lyophilized (freeze-dried) vaccine powder or wafer in one vial must be reconstituted (mixed) with the diluent (liquid) in another.

- Only use the diluent provided by the manufacturer for that vaccine as indicated on the chart.
- ALWAYS check the expiration date on the diluent and vaccine. NEVER use expired diluent or vaccine.

Vaccine product name	Manufacturer	Lyophilized vaccine (powder)	Liquid diluent (may contain vaccine)	Time allowed between reconstitution and use*	Diluent storage environment
ActHIB (Hib)	sanofi pasteur	Hib	0.4% sodium chloride	24 hrs	Refrigerator
Hiberix (Hib)	GlaxoSmithKline	Hib	0.9% sodium chloride	24 hrs	Refrigerator or room temp
Imovax (RAB _{HDCV})	sanofi pasteur	Rabies virus	Sterile water	Immediately [†]	Refrigerator
M-M-R II (MMR)	Merck	MMR	Sterile water	8 hrs	Refrigerator or room temp
MenHibrix (Hib-MenCY)	GlaxoSmithKline	Hib-MenCY	0.9% sodium chloride	Immediately [†]	Refrigerator or room temp
Menomune (MPSV4)	sanofi pasteur	MPSV4	Distilled water	30 min (single-dose vial) 35 days (multidose vial)	Refrigerator
Menveo (MCV4)	Novartis	MenA	MenCWY	8 hrs	Refrigerator
Pentacel (DTaP-IPV/Hib)	sanofi pasteur	Hib	DTaP-IPV	Immediately [†]	Refrigerator
ProQuad (MMRV)	Merck	MMRV	Sterile water	30 min	Refrigerator or room temp
RabAvert (RAB _{PCECV})	Novartis	Rabies virus	Sterile water	Immediately [†]	Refrigerator
Rotarix (RV1) [‡]	GlaxoSmithKline	RV1	Sterile water, calcium carbonate, and xanthan	24 hrs	Room temp
Varivax (VAR)	Merck	VAR	Sterile water	30 min	Refrigerator or room temp
YF-VAX (YF)	sanofi pasteur	YF	0.9% sodium chloride	60 min	Refrigerator or room temp
Zostavax (HZV)	Merck	HZV	Sterile water	30 min	Refrigerator or room temp

Always refer to package inserts for detailed instructions on reconstituting specific vaccines. In general, follow the steps below.

- For single-dose vaccine products (exceptions are Menomune in the multi-dose vial and Rotarix[‡]), select a syringe and a needle of proper length to be used for both reconstitution and administration of the vaccine. Following reconstitution, Menomune in a multidose vial will require a new needle and syringe for each dose of vaccine to be administered. For Rotarix, see the package insert.[‡]
- Before reconstituting, check labels on both the lyophilized vaccine vial and the diluent to verify that
 - they are the correct two products to mix together,
 - the diluent is the correct volume (especially for Menomune in the multidose vial), and
 - neither the vaccine nor the diluent has expired.
- Reconstitute (i.e., mix) vaccine **just prior to use** by
 - removing the protective caps and wiping each stopper with an alcohol swab,
 - inserting needle of syringe into diluent vial and withdrawing entire contents, and
 - injecting diluent into lyophilized vaccine vial and rotating or agitating to thoroughly dissolve the lyophilized powder.
- Check the appearance of the reconstituted vaccine.
 - Reconstituted vaccine may be used if the color and appearance match the description on the package insert.
 - If there is discoloration, extraneous particulate matter, obvious lack of resuspension, or the vaccine cannot be thoroughly mixed, mark the vial as “DO NOT USE,” return it to proper storage conditions, and contact your state or local health department immunization program or the vaccine manufacturer.
- If reconstituted vaccine is not used immediately or comes in a multidose vial (i.e., multi-dose Menomune), be sure to
 - clearly mark the vial with the date and time the vaccine was reconstituted,
 - maintain the product at 35°–46°F (2°–8°C); do not freeze, and
 - use only within the time indicated on chart above.

* If the reconstituted vaccine is not used within this time period, it must be discarded.

† For purposes of this guidance, “immediately” is defined as within 30 minutes or less.

‡ Rotarix vaccine is administered by mouth using the applicator that contains the diluent. It is not administered as an injection.